

Original researches

Effect of herbal feed additives on the quality of colostrum, immunological indicators of newborn calves blood and growth energy of young animals

V. S. Kozyr*, P. P. Antonenko**, R. V. Mylostyvyi**, N. I. Suslova**, P. M. Skliarov**, O. P. Reshetnychenko***, T. D. Pushkar***, V. O. Sapronova**, O. M. Pokhyl**

* Institute of Grain Crops of National Academy of Agrarian Sciences, Dnipro, Ukraine

** Dnipro State Agrarian and Economic University, Dnipro, Ukraine

*** Odessa State Agrarian University, Odessa, Ukraine

Received: 26 April 2019

Revised: 05 May 2019

Accepted: 21 July 2019

Institute of Grain Crops of National Academy of Agrarian Sciences, Volodymyr Vernadskyi Str., 14, Dnipro, 49027, Ukraine

Dnipro State Agrarian and Economic University, Sergii Efremov Str., 25, Dnipro, 49600, Ukraine

Odessa State Agrarian University, st. Panteleymonovskaya, 13, Odessa, 65012, Ukraine

Tel.: +38-097-280-88-19

E-mail: mylostyvyi.r.v@dsau.dp.ua

Cite this article: Kozyr, V. S., Antonenko, P. P., Mylostyvyi, R. V., Suslova, N. I., Skliarov, P. M., Reshetnychenko, O. P., Pushkar, T. D., Sapronova, V. O., & Pokhyl, O. M. (2019). Effect of herbal feed additives on the quality of colostrum, immunological indicators of newborn calves blood and growth energy of young animals. *Theoretical and Applied Veterinary Medicine*, 7(3), 137–142. doi: 10.32819/2019.71024

Abstract. Studies have shown that the use of herbal feed additives «Fitopank» by Grey Ukrainian cows 30 days before calving had a positive effect not only on the body of cows-mothers and the quality of their colostrum, but also on the physiological state of the born calves and the growth energy of young animals. Adding of «Fitopank» to the cows diet helped to improve the quality of colostrum and its immunological value. It showed a probable increase of total protein by 10–14 % ($P < 0.05$) and immunoglobulin level by 23–25 % ($P < 0.01$). The titratable acidity of both the first and second milk yields was 16 % lower ($P < 0.01$) than in the control group. The use of this feed additive improved the immunological status of newborn calves. In their blood there was an increase of total protein content by 17 % ($P < 0.05$), the level of gamma globulins by 18% ($p < 0.05$), IgG concentrations by 14 % ($p < 0.05$), Ig M by 13 % and Ig A by 17 % ($p < 0.05$), as well as the growth of T-lymphocytes by 46% and B-lymphocytes by 24 % ($p < 0.01$), compared with control group. Detected increased growth of the 8–12 months bulls (by 2.3%, $p < 0.05$) of the experimental group may indicate their tendency for better development of meat qualities. Experiment showed that the absolute live weight increase of young animals up to 18 months was higher than the control by 10.9 kg ($p < 0.05$). The study of the herbal supplements efficacy for the prevention and treatment of animals immunodeficiency, as well as their impact on the metabolic processes and productive qualities of farm animals will be the focus of our further research.

Keywords: beef cattle; feed additives; colostrum; immune status; growth energy.

Вплив рослинного препарату на якість молозива, імунологічні показники крові новонароджених телят та енергію їх росту

В.С. Козир*, П.П. Антоненко**, Р.В. Милостивий**, Н.І. Суслова**, П.М. Склярів**, О.П. Решетниченко***, Т.Д. Пушкар***, В.О. Сапронова**, О.М. Похил**

* Державна установа Інститут зернових культур НААН, Дніпро, Україна

** Дніпровський державний аграрно-економічний університет, Дніпро, Україна

*** Одеський державний аграрний університет, Одеса, Україна

Анотація. Встановлено, що застосування кормової фітодобавки «Фітопанк» коровам сірої української породи за 30 днів до отелення мало позитивний вплив не тільки на організм корів-матерів і якість їх молозива, а й на фізіологічний стан отриманих від них телят та енергію росту молодняка. Встановлено, що додавання до раціону «Фітопанку» сприяло підвищенню якості молозива та його імунологічної повноцінності. У ньому відбувалося вірогідне зростання загального білка – на 10–14 % ($P < 0,05$) та рівня імуноглобулінів – на 23–25 % ($P < 0,01$). Титрована кислотність як першого, так і другого удою була меншою на 16 % ($P < 0,01$), ніж у контролі. З'ясовано, що використання кормової фітодобавки покращувало імунологічний статус новонароджених телят. В їх крові відмічалось підвищення вмісту загального білка – на 17 % ($P < 0,05$), рівня гамма-глобулінів – на 18 % ($p < 0,01$), концентрації IgG – на 14% ($p < 0,01$), Ig M – на 13 % та Ig A – на 17 % ($p < 0,01$), а також зростання Т-лімфоцитів – на 46 % і В-лімфоцитів – на 24% ($p < 0,01$), порівняно з контролем. Виявлена перевага за відносною швидкістю росту в бугайців дослідної групи, порівняно з аналогами 8–12-місячного віку (на 2,3 %, $p < 0,05$), може свідчити про їх схильність до кращого формування м'ясних якостей. Абсолютний приріст живої маси молодняка до 18-місячного віку за впливу фітодобавки був вищим за контроль на 10,9 кг ($p < 0,01$). Вивчення ефективності препаратів (добавок) рослинного походження з метою попередження і лікування імунодефіцитних станів тварин, а також їх вплив на метаболічні процеси та продуктивні якості сільськогосподарських тварин будуть напрямом наших подальших досліджень.

Ключові слова: м'ясна худоба; кормова фітодобавка; молозиво; імунний статус; енергія росту.

Introduction

Newborn diseases are one of the most common diseases in animal husbandry. In particular, despite significant advances in growing and maintaining dairy calves in the United States, one in ten calves die before weaning (Hulbert & Moisa, 2016). For example, in the Netherlands (Santman-Berends et al., 2019), calf mortality before 1 year age is approximately 16.5 %. Dividing the first year of life into three age categories, the death rate for postnatal calves (≤ 14 days) was 3.3 %; early weaned (15 to 55 days) – 4.5% and weaned > 56 days – 3.1 %. Researchers (Voljč et al., 2019) report that about 8 % of prenatal calf deaths were related to the functional condition of cows, while only 4 % to neonatal pathology. In the latter case, the season of birth and the size of the herd had a significant impact. According to other scientists, the death rate of newborn calves was as high as 40 % (Kasap et al., 2019) and it was highest from birth to six months age, as it was associated not only with born calves but also with care and feeding of cows in the dry season. During this period cows may have a high negative energy balance and fatty liver, which affects the health of both mother and calf. The authors inform (Abuelo et al., 2019) that the increased risk of diseases in the prenatal and neonatal periods is a consequence of disturbance of the redox balance in the body and the manifestation of oxidative stress, so the use of additives containing natural antioxidants can prevent animal diseases.

The most important requirements for preventing neonatal pathology are care and feeding, and in particular, keeping the time and term of the first portion of colostrum, as its maternal antibodies are effective in preventing many diseases of newborn calves (Hulbert & Moisa, 2016). Insufficient amount of immunoglobulins synthesized by the fetus is not able to protect it effectively after birth. Immunoglobulins supplied with colostrum remain the only source of passive immunity, so the constant monitoring of its quality will allow reliable transmission of passive immunity to calves (Chernenko et al., 2017). Scientists believe that controlling of colostrum feeding and improving nutrition before heifers weaning will have a positive effect on the mammary gland formation (Vailati-Riboni et al., 2018) and further cow productivity (Makau et al., 2018; Milostiviy et al., 2018b), as immunobiological indicators of blood are closely linked with the productive qualities of livestock (Milostiviy et al., 2018a; Elsohaby et al., 2019).

It is very important to use the means for correcting the immunodeficiency states of productive and young animals. Despite the great arsenal of domestic and foreign medicines, they are not always effective and safe. Therefore, the use of biologically active natural additives of plant origin has become widespread in preventive veterinary practice, but their efficacy requires appropriate research. So, the purpose of the work was to determine the immunological value of cows colostrum, the immune status of newborn calves and the growth energy of young animals effecting by herbal feed supplement «Fitopank».

Material and methods

The study is a fragment of the research «Application of biologically active substances of plant origin for disease prevention,

Table 1. Composition of cow colostrum (n = 40, $\bar{X} \pm SE$)

Indicator	1-st milking		2-nd milking	
	control	experimental	control	experimental
Total protein, g / l	185.5 ± 6.46	204.6 ± 8.51*	165.0 ± 1.79	189.0 ± 2.03*
Immunoglobulins, g / l	48.2 ± 1.84	59.5 ± 2.25**	32.0 ± 1.07	40.1 ± 2.37**
Titrated acidity, °T	52.4 ± 1.41	44.0 ± 0.54**	42.3 ± 0.98	35.4 ± 1.02**

Note: * – p < 0.05, ** – p < 0.01 compared to control.

preservation and non-specific immunity» (state registration number 0112U08497) and «Impact of technological factors on the improving of the livestock products quality» (state registration number 0114U005590) Dnipro State Agrarian and Economic University. The experiment was performed on the Polivanivka cattle farm in Magdalynivka district, Dnipropetrovsk region. From the total herd of full-grown Grey Ukrainian cows in 30 days before calving, a control (n = 20) and experimental (n = 20) groups of cows-mothers and their calves (respectively n = 7 and n = 6) were formed on the principle of analogues (age, live weight, physiological state). Cows with calves were winter-stalled. All the animals involved in the experiment were clinically healthy. During the experiment, the requirements of the European Convention for the Protection of Vertebrate Animals Used for Experimental and Scientific Purposes (Strasbourg, 1986) and the Law of Ukraine “On the Protection of Animals against Cruelty” No. 3447 of 21.02.2006 were observed. The study material was serum and colostrum.

«Fitopank» feed additive is a complex composition of alcoholic infusions of seven medicinal plants combined in the author’s ratio (Filippov et al., 2002): the root of palmate rhubarb of Tungut (*Rheum palmatum* var. *Tanguticum*), dill (*Anethum graveans*, L.), root of wild sunflower (*Inula helenium* L.), the root of German iris (*Iris germanis* L.), poison Hemlock herb (*Conium maculatum* L.), the leaves of the bogbean (*Menyanthes trifoliata* L.) and sage (*Savlia officinalis*). «Fitopank» was given to cows individually at a dose of 1 ml twice a day for 30 days before calving with a rubber bottle, spreading it in 100 ml of water.

The number of erythrocytes and leukocytes in blood were counted by Goryaev’s method; hemoglobin was studied by hemoglobincyanide method; total protein, total calcium, inorganic phosphorus, reserve alkalinity in serum were determined by chemical analyzer «Stat Fox-1904» (USA). The level of gamma globulins was determined by the turbidimetric method an concentration of immunoglobulins A, G, M by radial immunodiffusion method of G. Mamcini et al., (1965). The number of T- and B-lymphocytes in the peripheral blood was determined by spontaneous rosette formation. Colostrum composition and its quality were investigated by conventional methods (Tsekhmistrenko & Kononskyi, 2014). Total collagen immunoglobulin content was determined by Kruise colostrometer (Denmark) using the appropriate scale on the instrument. Growth rates (average daily growth, relative live weight increase and live weight increase coefficient) for different age periods of the young animals were calculated on the basis of monthly weightings using the above methods (Pidpala, 2007).

The obtained data were statistically processed using the built-in mathematical functions of the Office Excel 2007 computer program and the Statistica for Windows 6.0 statistical suite with Student’s t test.

Results

The physiological status of the cows was examined before the application of «Fitopank» feed additive. It was established that the hematologic indices of cows in the control and experimental groups were within the reference values: the number of red blood cells were 5.6 ± 0.09 and 5.7 ± 0.07 T / l, the leukocyte number

Table 2. Immunological indices of ten-day-old calves due to the effects of «Fitopank», $\bar{X} \pm SE$

Indicator	Control group, n = 7	Experimental group, n = 6
Total protein, g / l	57.6 ± 2.54	67.4 ± 2.28*
Gamma globulins, g / l	17.0 ± 0.51	20.1 ± 0.75*
Immunoglobulins G, g / l	14.0 ± 0.37	16.0 ± 0.24*
Immunoglobulins M, g / l	1.4 ± 0.07	1.8 ± 0.11*
Immunoglobulins A, g / l	0.9 ± 0.12	1.5 ± 0.13*
T-lymphocytes, %	27.5 ± 1.41	40.2 ± 2.53**
B-lymphocytes, %	7.6 ± 0.26	9.4 ± 0.42**

Note: * – p < 0.05, ** – p < 0.01 compared to control.

was 9.1 ± 0, 16 and 9.0 ± 0.18 G/l, hemoglobin content was 05.4 ± 2.65 and 103.8 ± 3.96 g/l, total protein was 72.0 ± 1.58 and 71.8 ± 2.16 g/l, total calcium concentration was 2.59 ± 0.06 and 2.61 ± 0.11 mmol/l, inorganic phosphorus was 1.93 ± 0.05 and 2.01 ± 0.09mmol/l, the reserve alkalinity was 53.4 ± 0.96 and 52.7 ± 0.83 % CO² respectively. The differences we found between the groups of cows in terms of blood were unreliable and testify to the satisfactory condition of the animals in the dry season.

The use of feed herbal supplements affected the immunological value of cow colostrum. It was found (Table 1) that the content of immunoglobulins in colostrum of the second milk yield decreased by 34 % in the control group and 33 % in the experimental animals compared to the first milk (p < 0.001). However, due to the effect of «Fitopank» in the colostrum of both the first and the second milk, the level of immunoglobulins was higher by 23 and 25 %, than in the control (p < 0.01). Total protein content increased by 10 and 14%, (p < 0.05). The titratable acidity of colostrum was 16 % lower (p < 0.01) than in animals of the control group.

It was found (Table 2) that the application of «Fitopank» for cows before calving had a positive effect on the immunological status of calves body. In their blood there was an increase in total protein content by 17% (p < 0.05), the level of gamma globulins by 18 % (p < 0.05), Ig G concentrations by 14% (p < 0.05) , Ig M by 13 % and Ig A by 17 % (p < 0.05), as well as the growth of T-lymphocytes by 46 % and B-lymphocytes by 24 % (p < 0.01), compared with control group. All indicators were within the physiological norm.

Thus, biologically active substances in the composition of the herbal additives had a positive effect on the cows body, improving

the immunological status of calves in the postnatal period.

Live weight is one of the main economically useful traits of animal productivity in beef cattle that characterize growth, development and meat quality of young animals. The analysis of the obtained data (Table 3) shows that the bulls of the experimental groups grew intensively, developed well and in all ages according to the live weight met the requirements of the first class and elite. In the feeding period, differences in live weight of the control and experimental groups were minimal. However, with age, their growth pattern has changed. In particular, the live weight of the bulls in experimental group was by 8.3 and 6.9 kg heavier at 12 and 15 months age, and by 10.5 kg at 18 months (p < 0.05). Overall, the absolute increase in live weight of the experimental animals was higher than the control by 10.9 kg.

An important indicator characterizing the level of life productivity of young animals is the value of the live weight increase over certain periods of time. It should be noted that the average daily increase of experimental bulls during the whole period of rearing was high enough and the changes in keeping and feeding after weaning did not affect the health of the experimental young animals (Table 4).

It is significant that from 8 to 12 months, the average daily increase of experimental bulls was 767–830 gr, so, they effectively used herbal feed additives in this age period. However, the increase in live weight of the experimental group animals was higher than the control by 63.1 g or 8.2 % (p < 0.05). In the future (15–18 months) differences were less significant, but the predominance (7.3 %) of experimental bulls remained.

A more objective and complete dynamics of growth rate is

Table 3. Dynamics of live weight of small bulls, kg $\bar{X} \pm SE$

Age, month	Group	
	control, n = 7	experimental, n = 6
Newborns	28.7 ± 0.45	28.3 ± 0.44
6	183.6 ± 0.86	184.8 ± 1.02
8	230.5 ± 1.35	231.1 ± 1.69
12	324.1 ± 2.04	332.4 ± 3.54*
15	385.2 ± 2.18	392.1 ± 1.61*
18	436.2 ± 1.42	446.7 ± 2.04*
Absolute live weight increase	407.5 ± 1.98	418.4 ± 1.63*

Note: * – p < 0.05 compared to control.

Table 4. Average daily increase of bulls live weight, by periods of growth, g $\bar{X} \pm SE$

Age, month	Group	
	control, n = 7	experimental, n = 6
0–8	827.1 ± 5.99	831.1 ± 7.31
0–15	779.2 ± 2.97	795.2 ± 4.01
0–18	742.3 ± 2.93	762.1 ± 3.28
8–12	767.2 ± 13.62	830.3 ± 10.48*
8–15	724.6 ± 9.38	754.1 ± 9.91
8–18	674.4 ± 6.51	706.9 ± 6.67
12–18	612.6 ± 9.01	624.6 ± 9.96
15–18	557.4 ± 12.59	596.7 ± 15.13

Table 5. The relative growth rate and the coefficient of live weight increase of bulls, $\bar{X} \pm SE$

Indicator	Group	
	control, n = 7	experimental, n = 6
	Relative speed, %	
0–8 months	155.7 ± 0.66	156.4 ± 0.68
8–12 months	33.8 ± 0.63	36.1 ± 0.51*
8–15 months	50.3 ± 0.67	51.6 ± 0.71
8–18 months	61.7 ± 0.61	63.6 ± 0.69
0–15 months	172.3 ± 0.42	173.1 ± 0.44
0–18 months	175.3 ± 0.38	176.2 ± 0.37
	The coefficient of live weight increase	
6	6.41 ± 0.11	6.54 ± 0.11
8	8.05 ± 0.16	8.18 ± 0.14
12	11.31 ± 0.18	11.77 ± 0.21
15	13.45 ± 0.22	13.88 ± 0.24
18	15.22 ± 0.25	15.82 ± 0.26

Note: * – p < 0.05 compared to control.

given by the rate of relative growth rate and the rate of body weight increasing with age. It was found (Table 5) that the relative growth rate decreased with age and was almost at the same level. Youngsters of all groups grew and developed normally. The obtained data indicate a sufficiently high level of their productive qualities.

At the same time, better growth rate of bulls in the experimental group compared to control group (by 2.3 %, p < 0.05) at 8–12 months of age, may indicate better meat quality.

Discussion

Serum protein studies have been widely used to evaluate the functional status of newborn calves (Kalaeva et al., 2019). Usually determination of the concentration of total protein (STP) and immunoglobulin G (IgG) is a reliable method of determining the effectiveness of passive transfer of calves immunity (Wilm et al., 2018; Gavrilin et al., 2018). However, data on their normal serum content of newborn calves are quite limited, especially in the early days of the postnatal period. Scientists report (Staněk et al., 2019) that the concentration of IgG determined in the serum of newborn calves by radial immunodiffusion averaged 13.7 g/l (with fluctuations in the range of 1.5 to 46.6 g/l) and significantly depended on the breed, herd size and period of year. Guided by the criterion of IgG < 10 g/l, scientists have concluded that 34.6 % of calves had passive transfer immunity syndrome (FPT).

The concentration of IgG in the serum of newborns is an important factor in preventing the morbidity and death of calves (Elsobaby et al., 2019) and depends on the quality of colostrum. Citing data from foreign researchers, it is reported (Chernenko et al., 2017) that mortality among calves with serum Ig less than 12 mg/ml 24.8 hours after birth was 6.8 %, and with a blood Ig concentration above 12 mg / ml only 3.3 %. It was noted (Wilm et al., 2018) that immediately after birth, before feeding the colostrum, the content of STP and IgG were respectively at the levels of 46.1 and 6.0 g/l. After 24 h after feeding the first portions of colostrum, their concentration was at the level of 58.3 and 22.2 g/l, and on the tenth day it was 57.8 and 16.1 g/l, respectively. Therefore, IgG concentration, compared to its maximum initial serum level (24 h after the first colostrum intake), «decreased» at a rate of 0.69 g/l every day. Using a total protein of < 55 and 60 g/l as a passive

immunity failure assessment, researchers (Pempek et al., 2017) found that 7,6–26,8 % of all auctioned calves in Ohio (USA) had it. 18.8 % of calves had clinical signs of diarrhea, 19.3 % had dyspepsia and 30.7 % had navel inflammation. But only 0.5 % (0 to 1.3 %) of calves had respiratory diseases.

It was noted (Ellis et al., 2018) that feeding colostrum with 150g of IgG provided sufficient production of antibodies to respiratory syncytial virus (PC virus) in the nasal mucosa of calves, and therefore the study of colostrum in a practical environment using a special hydrometer (colostrometer) can be considered as one of the simple and reliable methods of quality control (Chernenko et al., 2017). The existence of a reliable relationship between the content of immunoglobulins of classes G and M in the serum of three-month-old calves with signs of their milk productivity (r = 0,31-0,63; P < 0,05), allows to confirm the feasibility of immunological studies of blood for early prognosis of dairy cow productivity (Milostiviy et al., 2018a).

It has been reported (Abuelo et al., 2019) that the colostrum content of immunoglobulins of class G and total protein depended on stress resistance of cows, and indirectly affected the viability of their calves. So, increase of total protein and immunoglobulins in the colostrum of cows by application of the biologically active substances of plant components of «Fitopank», will positively affect the immunological parameters of newborn calves, which is proven by our studies.

It is known (Hulbert & Moisés, 2016) that behavioral responses and stress of calves after weaning may affect their immunity when transferred from individual to group retention. Various stressors at an early age (decoration, castration, etc.) lead to a decrease in passive transmission of immunoglobulins and the development of immunodeficiency. Despite the availability of anesthetics and analgesics, the evidence suggests that they also cause leukocyte function inhibition when used during these painful procedures. Therefore, the use of natural herbal additives for the prevention and correction of immunodeficiency of animals is a promising research area that deserves attention.

Unfortunately, there is limited data in the scientific literature regarding the mechanism of action of many herbal immunotropic drugs. Their immunomodulatory effects are usually associated with the production of cytokines and mediators (Markova, 2004). This is quite justified, as the immune system works by the interaction

of T-, B-cells, natural killer cells (EC cells), macrophages and mediators (interferons, interleukins, etc.). We are inclined to believe that the improvement of colostrum quality and, as a result of the immunological status of newborn calves due to the effects of «Fitopank», are directly related to the composition of the plants that are part of this feed additive. In particular, about 200 compounds are isolated from rhubarb, including anthraquinones, anthrons, stilbenes, flavonoids, acylglycosides and nails. These components have great pharmacological activity, including antitumor, hepatitis, anti-inflammatory, antimicrobial and analgesic action (Cao et al., 2017). Dill essential oils have antibacterial, antifungal, antioxidant, insecticidal, anti-inflammatory, antispasmodic and others properties (Chahal et al., 2017). Numerous types of wild sunflower are used alone or as an important ingredient in various drug for the treatment of cardiovascular, respiratory, urinary, central nervous and digestive disorders, for the treatment of skin, liver, fungal and bacterial infections. A number of phytochemicals, including alkaloids, essential oils, flavonoids, terpenes and lactones isolated from herbs of the *Inula* sorts, explain the traditional use of these plants as medicines (Das et al., 2019). Types of iris are well known as sources rich in isoflavonoids, which have antioxidant and antidiabetic activity (Ibrahim et al., 2017), capable of exhibiting anti-ulcer and anti-inflammatory properties, having a positive effect in chronic neurodegenerative processes (Borhani et al., 2017). Bogbean is rich in chemicals (betulinic acids and triterpene saponins) with anti-inflammatory and immunomodulatory effects, and also has antineoplastic properties associated with the induction of apoptosis in neoplastic cells (Kowalczyk et al., 2019). Polyphenols and biologically active compounds in the composition of sage have a therapeutic effect in neurodegenerative diseases and various cancers (Jakovljević et al. 2019). However, some components of «Fitopank» have strong toxic properties. In particular, hemlock causes an intoxication with the hypersalivation, anorexia, hyperthermia, tachycardia, hyperpnea, muscle spasms, unsteady gait, ataxia and hyperreflexia, and even death (Cortinovic & Caloni, 2015). Therefore, the use of herbal feed additives to correct the immune deficiency of the body is a promising research direction. However, the ratio and content of the active components of the plants should be a subject of careful study, as some of them may have strong toxic properties.

Conclusions

The effectiveness of herbal feed additives for the prevention and treatment of animals immunodeficiency is very promising scientific research in the field of veterinary medicine. Application of the herbal feed additives «Fitopank» for dry cows helped to increase the level of immunoglobulins in colostrum by 23–25 % ($P < 0.01$), increase the level of immunoglobulins by 13–17 % ($P < 0.05$) in serum of newborn calves and T- and B lymphocytes by 24–46 % ($p < 0.01$), as well as an increase of relative growth rate of bulls and their absolute growth to 18 months age.

References

Abuelo, A., Hernández, J., Benedito, J., & Castillo, C. (2019). Redox Biology in Transition Periods of Dairy Cattle: Role in the Health of Periparturient and Neonatal Animals. *Antioxidants*, 8(1), 20.

Borhani, M., Sharifzadeh, M., Farzaei, M. H., Narimani, Z., Sabbaghziarani, F., Gholami, M., & Rahimi, R. (2017). Protective effect of *Iris germanica* L. in β -amyloid-induced animal model of alzheimer's disease. *African Journal of Traditional, Complementary and Alternative Medicines*, 14(4), 140–148.

Cao, Y.-J., Pu, Z.-J., Tang, Y.-P., Shen, J., Chen, Y.-Y., Kang, A., Zhou, G.-S., Duan, J.-A. (2017). Advances in bio-active constituents, pharmacology and clinical applications of rhubarb.

Chinese Medicine, 12(1).

Chahal, K.K., Monika, Kumar, A., Bhardwaj, U., & Kaur, R. (2017). Chemistry and biological activities of *Anethum graveolens* L. (dill) essential oil: A review. *Journal of Pharmacognosy and Phytochemistry*, 6(2), 295–306.

Chernenko, O. M., Chernenko, O. I., & Sanjara, R. A. (2017). The quality of colostrum and vitality of calves, born from cows with different reaction to stress experiences. *Regulatory Mechanisms in Biosystems*, 8(2), 299–303.

Cortinovic, C., & Caloni, F. (2015). Alkaloid-containing plants poisonous to cattle and horses in Europe. *Toxins*, 7(12), 5301–5307.

Das, A., Shakya, A., Ghosh, S.K., Singh, U.P., & Bhat H.R. (2019). A review of phytochemical and pharmacological studies of *Inula* species. *Current Bioactive Compounds*.

Ellis, J.A., Chamorro, M.F., Lacoste, S., Gow, S.P., & Haines, D.M. (2018). Bovine respiratory syncytial virus-specific IgG-1 in nasal secretions of colostrum-fed neonatal calves. *The Canadian Veterinary Journal*, 59(5), 505–508.

Elsohaby I., Cameron M., Elmoslemany A., McClure J. T., & Keefe G. (2019). Effect of passive transfer of immunity on growth performance of preweaned dairy calves. *Canadian Journal of Veterinary Research*, 83(2), 90–96.

Filippov, Yu.O., Antonenko, P.P., & Ardelian, V.M. (2002). Patent na korysnu model № 47289, 6A61K35/78 «Sposib korektsii imunnoi systemy u molodniaka silskohospodarskykh tvaryn» [A method for correcting the immune system in young animals of farm animals]. Zaiavl. 16.10.2001. Opubl. 17.06.2002. Biul. №6 (in Ukrainian).

Gavrilin, P. N., Lieshchova, M. A., Gavrilina, O. G., & Boldyreva, T. F. (2018). Prenatal morphogenesis of compartments of the parenchyma of the lymph nodes of domestic cattle (*Bos taurus*). *Regulatory Mechanisms in Biosystems*, 9(1), 95–104.

Hulbert, L. E., & Moisés, S. J. (2016). Stress, immunity, and the management of calves. *Journal of Dairy Science*, 99(4), 3199–3216.

Ibrahim, S. R. M., Mohamed, G. A., Zayed, M. F., & Ross, S. A. (2017). 8-Hydroxyirilone 5-methyl ether and 8-hydroxyirilone, new antioxidant and α -amylase inhibitors isoflavonoids from *Iris germanica* rhizomes. *Bioorganic Chemistry*, 70, 192–198.

Jakovljević, M., Jokić, S., Molnar, M., Jašić, M., Babić, J., Jukić, H., & Banjari, I. (2019). Bioactive profile of various *salvia officinalis* L. Preparations. *Plants*, 8(3), 55.

Kalaeva, E., Zemlyanukhina, O., Kalaev, V., Efimova, K., Chernitskiy, A., Kaverin, N., & Safonov, V. (2019). The influence of haptoglobin phenotype on differential leukocyte count in neonatal calves. *Turkish Journal of Veterinary and Animal Sciences*, 43(2), 177–185.

Kasap, S., Temizel, E. M., Akgul, G., & Senturk, S. (2018). 0-6 Aylık buzağlarda bulaşıcı hastalıkların azaltılması ve sonuçları için pratik saha uygulamaları. *Harran Üniversitesi Veteriner Fakültesi Dergisi*, 7(1), 102–107.

Kowalczyk, T., Sitarek, P., Skala, E., Toma, M., Wielanek, M., Pytel, D., Wiczfińska, J., Szemraj, J., and Śliwiński, T. (2019). Induction of apoptosis by in vitro and in vivo plant extracts derived from *Menyanthes trifoliata* L. in human cancer cells. *Cytotechnology*, 71(1), 165–180.

Makau, D. N., VanLeeuwen, J. A., Gitau, G. K., Muraya, J., McKenna, S. L., Walton, C., & Wichtel, J. J. (2018). Animal and management factors associated with weight gain in dairy calves and heifers on smallholder dairy farms in Kenya. *Preventive Veterinary Medicine*, 161, 60–68.

Markova, T.P. (2004). Immunotropnyie preparaty. Za i protiv [Immunotropic drugs. Pros and cons]. *Remedium*, 11, 45–49 (in Russian).

Milostiviy, R., Antonenko, P., Kostyuk, V., Vasilenko, T., & Czerniawska-Piątkowska, E. (2018a). The milk yield of Ukrainian Holstein is related to the immunobiological parameters of blood

- of calves. *Folia Pomeranae Universitatis Technologiae Stetinensis Agricultura, Alimentaria, Piscaria et Zootechnica*, 345(48), 163–172.
- Milostiviy, R.V., Nadraha, M.M., & Novokshonova, H.K. (2018b). Zviazok mizh udoiem koriv i yikh rostom v ontogenezi za riznoi tryvalosti molochnoho periodu [The relationship between milk yield of cows and their growth in ontogenesis depends on the duration of the period of milk intake]. *Visnyk Dnipropetrovskoho Derzhavnogo Ahrarno-Ekonomichnoho Universytetu*, 1–2(47), 29–35 (in Ukrainian).
- Pempek, J., Trearchis, D., Masterson, M., Habing, G., & Proudfoot, K. (2017). Veal calf health on the day of arrival at growers in Ohio. *Journal of Animal Science*, 95(9), 3863.
- Pidpala, T.V. (2007) Skotarstvo i tekhnolohiia vyrobnytstva moloka ta yalovychny [Livestock and technology for the production of milk and beef]. *Vydavnychi viddil MDAU. Mykolaiv* (in Ukrainian).
- Santman-Berends, I. M. G. A., Schukken, Y. H., & van Schaik, G. (2019). Quantifying calf mortality on dairy farms: Challenges and solutions. *Journal of Dairy Science*.
- Staněk, S., Nejedlá, E., Fleischer, P., Pechová, A., & Šlosárková, S. (2019). Prevalence of Failure of Passive Transfer of Immunity in Dairy Calves in the Czech Republic. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 67(1), 163–172.
- Tsekhmistrenko, S.I., & Kononskyi, O.I. (2014). *Biokhimiia molo-ka i molokoproduktiv* [Biochemistry of milk and milk products]. Bila Tserkva, (in Ukrainian).
- Vailati-Riboni, M., Bucktrout, R. E., Zhan, S., Geiger, A., McCann, J. C., Akers, R. M., & Looor, J. J. (2018). Higher plane of nutrition pre-weaning enhances Holstein calf mammary gland development through alterations in the parenchyma and fat pad transcriptome. *BMC Genomics*, 19(1).
- Voljč, M., Čepon, M., Malovrh, Š., & Žgur, S. (2019). Risk factors and mortality rate of calves in the first month of life in Slovenian Holstein Friesian population. *Journal of Central European Agriculture*, 20(1), 25–30.
- Wilm, J., Costa, J. H. C., Neave, H. W., Weary, D. M., & von Keyserlingk, M. A. G. (2018). Technical note: Serum total protein and immunoglobulin G concentrations in neonatal dairy calves over the first 10 days of age. *Journal of Dairy Science*, 101(7), 6430–6436.