

UNIVERSITY OF AGRONOMIC SCIENCES
AND VETERINARY MEDICINE OF BUCHAREST
FACULTY OF ANIMAL PRODUCTIONS
ENGINEERING AND MANAGEMENT

SCIENTIFIC PAPERS
SERIES D
ANIMAL SCIENCE
VOLUME LXVI, No. 2

2023
BUCHAREST

SCIENTIFIC COMMITTEE

- Stelian ACATINCĂI - University of Life Sciences "King Mihai I" from Timișoara, Romania
- Vasco A.P. CADAVEZ - Departamento de Ciência Animal & Centro de Investigação de Montanha (CIMO), Escola Superior Agrária, Instituto Politécnico de Bragança, Portugal
- Mioara COSTACHE - S.C.D.P. Nucet
- Cătălin DRAGOMIR - National Research-Development Institute for Animal Biology and Nutrition Balotești, Romania
- Nicolae EREMIĂ - State Agrarian University of Moldova, Chișinău, Republic of Moldova
- Horia GROȘU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Khalid Hamid HASSAN - University of Diyala, Irak
- Armagan HAYIRLI - Ataturk University, Erzurum, Turkey
- Mostafa A.R. IBRAHIM - University of Kafrelsheikh, Egipt
- Ondrej KADLECÍK - Slovak Agricultural University Nitra, Slovakia
- Yusuf KONCA - Erciyes University, Kayseri, Turkey
- Monica Paula MARIN - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Carmen Georgeta NICOLAE - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mircea POP - "Ion Ionescu de la Brad" University of Life Sciences Iasi, Romania
- Agatha POPESCU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Elena POPESCU-MICLOȘANU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Paul Rodian TĂPĂLOAGĂ - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Ilie VAN - Academy of Agricultural and Forestry Sciences "Gheorghe Ionescu-Șișești" Bucharest, Romania
- Livia VIDU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania

EDITORIAL BOARD

General Editor: Prof. Ph.D. Gheorghe Emil MĂRGINEAN

Executive Editor: Prof. Ph.D. Monica Paula MARIN

PUBLISHERS:

University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania – Faculty of Animal Science Engineering and Management

Address: 59 Mărăști Blvd, District 1, 011464, Bucharest, Romania

Phone: + 40 213 182 564, Fax: +40 213 182 888, www.zootehnie.ro

CERES Publishing House

Address: 29 Oastei Street, District 1, Bucharest, Romania

Phone: + 40 317 90 23, E-mail: edituraceres@yahoo.com, Webpage: www.editura-ceres.ro

Copyright 2023

To be cited: Scientific Papers, Series D. Animal Science, Volume LXVI, No. 2, 2023

*The publishers are not responsible for the opinions published in the Volume.
They represent the authors' point of view.*

ISSN 2285-5750; ISSN CD-ROM 2285-5769; ISSN Online 2393-2260; ISSN-L 2285-5750

International Database Indexing: Web of Science Core Collection (Emerging Sources Citation Index), Index Copernicus, CABI, DOAJ, Ulrich's Periodicals Directory (ProQuest), PBN, Cite Factor (Academic Scientific Journals), Scipio, OCLC (WorldCat), Research Bible, Google Scholar.

4. STUDY OF THE IMPACT OF SOME FACTORS ON THE GESTATION LENGTH OF ANGLO-NUBIAN GOATS REARED IN FOOTHILLS - Lora MONDESHKA, Svetoslava STOYCHEVA, Nikolay MARKOV, Miroslav HRISTOV, Tsvetelina DIMITROVA	244
5. THE COMPOSITION OF HEAVY METALS AND THE CONTENT OF ESTERIFIED FATTY ACIDS IN BEE TISSUES DEPENDING ON THE ENVIRONMENTAL CONDITION - Yosyp RIVIS, Volodymyr POSTOYENKO, Oleg STASIV, Olga STADNYTSKA, Bogdan GUTYJ, Svetlana USENKO, Bogdan SHAFERIVSKYI, Tetyna KARUNNA, Olena BEZALTYCHNA, Valentina YASKO, Olga HOPANENKO, Oleksandr DIACHENKO, Oleg KLUM, Ivan SARANCHUK	250
6. THE INFLUENCE OF POLYPHENOLS OF GREEN WALNUT EXTRACT ON ZINC HOMEOSTASIS AND ITS ROLE IN THE ORGANISM OF BREEDING ROOSTERS - Nicolae ROȘCA, Ion BALAN, Vladimir BUZAN, Sergiu BALACCI, Ion MEREUȚA, Iulia CAZACOV, Vasile HAREA, Melania BUCARCIUC, Vlad TEMCIUC, Ecaterina VÎHRIST	262
7. IMMUNOGLOBULINE-G LEVEL AND BODY WEIGHT OF BALB/C RECEIVING COMBINATION TREATMENT OF LYOPHILIZED <i>Curcuma longa</i> AND <i>Curcuma xanthorrhiza</i> DURING AN ANGIOGENESIS EXPERIMENTAL - Laurentius RUMOKOY, Jimmy POSANGI, Nontje Juliana KUMAJAS, Wisje Lusia TOAR	269

IV. SESSION TECHNOLOGIES OF ANIMAL HUSBANDRY

1. RESEARCH ON EFFECT OF MILKINGS FREQUENCY ON COWS' MILK PRODUCTION - Dumitru BACALU, Paula POSAN, Monica Paula MARIN, Alina UDROIU, Raluca Ioana PASCU, Carmen Georgeta NICOLAE	277
2. INVESTIGATIONS CONCERNING THE EXCRETION OF ANTIBIOTIC RESIDUES IN THE MILK OF COWS TREATED WITH ANTIBIOTICS - Mugurel COLA, Florica COLA	283
3. OPTIMIZATION OF INDOOR MICROCLIMATE PARAMETERS IS AN IMPORTANT FACTOR IN STIMULATING METABOLISM IN THE BODY AND INCREASING PIG PRODUCTIVITY - Andriyana DMYTROTSA, Stakh VOVK, Maria VOROBEL, Oleg KLYM	289
4. STUDY REGARDING THE IMPROVEMENT OF MILK PRODUCTION ACCORDING TO THE SIRE VALUES - Dănuț Nicolae ENEA, Sonia BEN FRAJ, Carmen Georgeta NICOLAE, Gheorghe Emil MĂRGINEAN, Livia VIDU	294
5. AI BASED DEVELOPMENT OF A LOW COMPUTATIONAL INTENSITY ALGORITHM FOR CATTLE HEART RATE (HR) ESTIMATION - Eva HAJNAL	300
6. LIMOUSIN BREED – CREATION, APPROVAL, SPECIFICATIONS AND CHALLENGES. REVIEW - Miroslav HRISTOV, Nikolay MARKOV, Tsvetelina DIMITROVA, Lora MONDESHKA, Svetoslava STOYCHEVA	308
7. INFLUENCE OF THE TEMPERATURE-HUMIDITY INDEX ON SOME PHYSIOLOGICAL PARAMETERS IN DAIRY GOATS - Romyana IVANOVA, Hristo HRISTEV	316
8. DISTRIBUTION AND PRODUCTIVE CHARACTERISTICS OF NORMANDE CATTLE BREED WORLDWIDE, IN EUROPE AND BULGARIA - REVIEW - Tatyana IVANOVA, Tsvetan MARKOV, Lora MONDESHKA, Miroslav HRISTOV	323
9. RESEARCH ON THE ECONOMIC ADVANTAGES OF BREEDING AUBRAC BEEF CATTLE: A REVIEW - Bianca-Maria MADESCU, Roxana LAZAR, Andrei Cristian MATEI, Paul Corneliu BOISTEANU	331
10. THE ANALYSIS OF DAIRY COWS FARM USING INTEGRATIVE NUMERICAL SYSTEM ANI 35L/2000 - George NICA, Bianca GROSEANU, Livia VIDU	342

THE COMPOSITION OF HEAVY METALS AND THE CONTENT OF ESTERIFIED FATTY ACIDS IN BEE TISSUES DEPENDING ON THE ENVIRONMENTAL CONDITION

Yosyp RIVIS¹, Volodymyr POSTOYENKO², Oleg STASIV¹, Olga STADNYTSKA¹, Bogdan GUTYJ³,
Svetlana USENKO⁴, Bogdan SHAFERIVSKYI⁴, Tetyna KARUNNA⁴, Olena BEZALTYCHNA⁵,
Valentina YASKO⁵, Olga HOPANENKO⁶, Oleksandr DIACHENKO¹, Oleg KLUM¹,
Ivan SARANCHUK⁷

¹Institute of Agriculture of the Carpathian Region of NAAS of Ukraine, Obroshino,
5 Hrushevskoho St., Lviv region, 81100, Ukraine

²Institute of Beekeeping named after P. I. Prokopovich, 19 Zabolotny st., Kyiv, 03680, Ukraine

³Stepan Gzhytskyi National University of Veterinary Medicine and Biotechnologies Lviv,
50 Pekarska st., Lviv, 79010, Ukraine

⁴Poltava State Agrarian University, 1/3 Skovorody st., Poltava, 36006, Ukraine

⁵Odessa State Agrarian University, 13 Panteleimonivska st., Odessa, 65012, Ukraine

⁶Higher Educational Communal Institution of Lviv Regional Council "Andrey Krupynsky Lviv
Medical Academy", 70 P. Doroshenko st., Lviv, 79000, Ukraine

⁷Bukovyna State Agricultural Experimental Station of NAAS, 21-A Bohdan Kryzhanovskiy St.,
Chernivtsi, 58026, Ukraine

Corresponding author email: bvh@ukr.net

Abstract

The work aimed to record the content of heavy metals, including toxic and esterified forms of fatty acids in tissues, and the honey productivity of bees in different natural zones of the Carpathian region. The level of dangerous elements of the first class of toxicity - Lead and Cadmium - increases significantly in the tissues of honey bees of the foothills and forest-steppe zones, compared to the conditionally clean mountain environment. In the Carpathian region, all heavy metals accumulate in more significant quantities in the abdominal tissues of honey bees than in the tissues of the chest and head. In the direction from the mountain to the foothills and further to the forest-steppe zone of the Carpathian region, a decrease in the content of esterified forms of fatty acids is observed in the tissues of honey bees. The intensity of conversion of the esterified form of linolenic acid into its longer-chain and unsaturated fatty acids of the omega-3 family in the tissues of the head of honeybees in the foothills and especially in the forest-steppe zones is sharply reduced.

Key words: *biomindicator, fatty acids, heavy metals, tissues.*

INTRODUCTION

The sources of heavy metal emissions and the ways of their entry into the environment are diverse. Still, they are generally of artificial origin due to urbanization and industrialization (Slivinska et al., 2018; 2019; 2020; 2021). Urbanization and industrialization, in particular, the activities of industry, agriculture, energy, and transport, as well as intensive extraction of minerals – all led to the release of heavy, including highly toxic, lead, cadmium, and arsenic into the air, water, soil and plants (Gutyj et al., 2017; 2019; Kovka & Nedashkivskiy, 2019; Tikhonova et al., 2020; Briffa et al., 2020).

The migration of heavy metals in objects of the external environment caused their accumulation in soils and plants (Krainiukov et al., 2020; Tikhonova et al., 2020; Briffa et al., 2020; Bashchenko et al., 2020; Prychepa et al., 2021). As a result, some types of plants gave way to others, and the terms of their flowering changed, thus the conditions of honey collection by bees (Kryvyi et al., 2018; Kovka & Nedashkivskiy, 2019; Krainiukov et al., 2020; Kobysch et al., 2021; Al-Kahtani et al., 2021). The above also led to the accumulation of heavy metals in the tissues of honey bees and beekeeping products (Didaras et al., 2020; Mărgăoan et al., 2021; Ricigliano et al., 2021).

As is known, heavy metals, including toxic ones, are involved in synthesizing, desaturation, and peroxide oxidation of long-chain fatty acids in plant and animal tissues and fluids. As a result, the supply of plant tissues, in particular pollen and bees, with energetic, structural, biologically active, and antimicrobial material changes (Desbois & Smith, 2010; Giri et al., 2018; Corby-Harris et al., 2021; Butsiak et al., 2021). All this affects bee colonies' productivity and their products' quality indicators (Matin et al., 2016; Kovalskiy et al., 2018; Vishchur et al., 2019; Gizaw et al., 2020; Monchanin et al., 2021).

Therefore, the issue of producing ecologically safe beekeeping products is urgent. Moreover, the production of honey bees occupies a prominent place in human life (Matin et al., 2016; Didaras et al., 2020; Costa et al., 2021; Ricigliano et al., 2021; El-Seedi et al., 2022). Very high requirements are placed on the quality indicators of honey bee products because Ukraine has become the leading exporter of honey to Europe.

In the literature, there needs to be more data on the content of heavy metals and different forms of fatty acids in bee pollen and tissues of honey bees kept in different natural zones of the Carpathian region (Klym & Stadnytska, 2019; Saranchuk et al., 2021).

Given the above, studies of the content of heavy metals, including toxic and esterified forms of fatty acids in tissues, and honey productivity of worker bees in different natural zones of the Carpathian region are of significant scientific and practical interest.

The goal of the work. To record the content of heavy metals, including toxic and esterified forms of fatty acids in tissues and honey productivity of bees in different natural zones of the Carpathian region.

MATERIALS AND METHODS

Experimental apiaries of clinically healthy honey bees of the Carpathian breed (*Apis mellifera* (L.) *carpatica*) were selected based on the private mountain (Slavsko village, Stryi district), foothills (Nizhnya Stynava village, Stryi district) and forest-steppe (village Myklashiv, Lviv district) zones of Lviv region,

where natural and climatic conditions and ecological situation are different.

To assess the intensity of technogenic load on the environment where experimental honey bee apiaries are located, the content of heavy metals (Ferrum, Zinc, Copper, Chromium, Cobalt, Nickel, Lead, and Cadmium) in the topsoil, bee pollen, and tissues of honey bees was determined.

In each of the above-described natural zones of the Carpathian region, the honey productivity of worker bees was studied in 3 apiaries and each of 5 hives (Kovalskiy & Kyryliv, 2011). In particular, the honey productivity of bees was reviewed in early June, mid-July, and mid-August by pumping out honey frames. These studies were conducted based on one bee colony of average strength and the same queen age. In addition, samples of bee pollen and honey bees were taken for laboratory research at each apiary from 3 hives in the spring-summer period. At the same time, examples of the arable layer of the soil were born in the radius of the available flight of honey bees.

The content of heavy metals, including toxic ones, was determined in selected samples of the topsoil, bee pollen, and tissues of the abdomen, chest, and head of honeybees and the pieces of the tissues above of honeybees – esterified forms of fatty acids. At the same time, the value of the level of heavy metals, including toxic and esterified forms of fatty acids, in tissues for the body of honey bees in different natural zones of the Carpathian region was analyzed.

The content of heavy metals (Ferrum, Zinc, Copper, Cobalt, Chromium, Nicol, Lead, and Cadmium) in the selected samples of the arable layer of the soil, bee comb, and tissues of the abdomen, chest, and head of honey bees was determined according to the currently valid state standard (DSTU 4405: 2005) on an atomic absorption spectrophotometer - Selma-115 (Vlizlo, 2012). The content of esterified forms of fatty acids in the tissues mentioned above of honey bees was determined by gas-liquid chromatography (Rivis et al., 2022).

The obtained digital material was processed by the method of variational statistics using the Student's criterion (Ibatullin & Zhukorskyi, 2017).

Arithmetic mean values (M) and errors ($\pm m$) were calculated. Differences were considered probable at $p < 0.05$. The computer program Origin 6.0 and Microsoft Excel were used for calculations.

RESULTS AND DISCUSSIONS

It was established that in the arable soil layer of the foothills and forest-steppe zones of the Carpathian region, compared to the

conditionally clean mountain zone, there is probably a higher content of the studied heavy metals (Table 1). At the same time, the arable layer of the soil of the forest-steppe zone of the Carpathian region contains the highest level of the studied heavy metals. The content of hazardous elements of the first toxicity class - Lead and Cadmium - in the arable layer of the soil in the above zone is 1.1 times higher than the maximum permissible concentration (Yatsuk & Balyuk, 2019).

Table 1. Gross content of heavy metals, including toxic ones, in the topsoil in different natural zones of the Carpathian region, g·10⁻³/kg of air-dry mass (M \pm m, n = 3)

Metal and its symbol	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Ferrum, Fe	14325.00 \pm 294.214	15184.29 \pm 454.862*	16573.04 \pm 294.429**
Zinc, Zn	47.58 \pm 4.488	78.52 \pm 3.722**	96.13 \pm 4.890***
Cuprum, Cu	21.60 \pm 1.391	34.56 \pm 1.828**	45.64 \pm 2.264***
Cobalt, Co	11.76 \pm 0.375	13.63 \pm 0.560**	17.20 \pm 1.830***
Chrome, Cr	41.69 \pm 2.283	63.65 \pm 3.584**	87.53 \pm 4.163***
Nicole, Ni	21.24 \pm 1.625	41.33 \pm 2.512***	59.42 \pm 3.214***
Lead, Pb	19.37 \pm 0.784	25.83 \pm 1.442*	33.30 \pm 2.870***
Cadmium, Cd	2.03 \pm 0.088	2.60 \pm 0.115*	3.20 \pm 0.271***

Note. Here and further, differences are likely compared to the mountainous zone: * - $p < 0.05$ -0.02; ** - $p < 0.01$; *** - $p < 0.001$

It is believed that the increase in the content of lead in the arable layer of the soil is associated with the intensive movement of motor vehicles (Loretta et al., 2015) and cadmium - with the introduction of meliorants and mineral fertilizers, primarily phosphogypsum and superphosphate, respectively (Razanov et al., 2015; Vozhehova et al., 2021). It can be seen that phosphoric acid residues in phosphogypsum and superphosphate bind solid elements and carriers for cadmium.

The obtained data characterize artificial environmental pollution in the experimental territories. The high level of heavy metals, including toxic ones, in the soil, is the reason for the increase in their concentration in bee pollen (plant pollen) obtained in the foothills and forest-steppe zones of the Carpathian region (Table 2). All this is a consequence of greater urbanization and industrialization of the above territories.

Table 2. The content of heavy metals, including toxic ones, in bee pollen in different natural zones of the Carpathian region, g·10⁻³/kg air-dry mass (M \pm m, n = 3)

Metal and its symbol	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Ferrum, Fe	33.52 \pm 0.830	37.11 \pm 0.781*	43.39 \pm 2.253**
Zinc, Zn	34.39 \pm 1.91	39.20 \pm 0.900*	42.72 \pm 0.872**
Cuprum, Cu	2.01 \pm 0.089	3.02 \pm 0.169*	4.20 \pm 0.170***
Cobalt, Co	1.01 \pm 0.029	1.14 \pm 0.050*	1.44 \pm 0.112***
Chrome, Cr	4.10 \pm 0.177	5.02 \pm 0.180*	6.68 \pm 0.149***
Nicole, Ni	0.58 \pm 0.015	0.65 \pm 0.015*	0.74 \pm 0.023**
Lead, Pb	0.12 \pm 0.007	0.18 \pm 0.009*	0.26 \pm 0.012**
Cadmium, Cd	0.04 \pm 0.003	0.07 \pm 0.007*	0.10 \pm 0.009**

It should be noted that in the forest-steppe zone of the Carpathian region, compared to the mountain zone, the arable layer of the soil and bee pollen has a relatively high content of probiotic heavy metals - Zinc, Copper, Cobalt, Chromium, and Nickel. The

above-mentioned heavy metals in acceptable quantities are essential for plant tissues and bees (Hsu et al., 2021). However, the increased level of toxic lead and cadmium in the arable layer of the soil and bee hives can neutralize the positive effect

of probiotic heavy metals on the mentioned tissues (Purać et al., 2019).

The high level of Ferrum, Zinc, Copper, Cobalt, Chromium, Nickel, Plumbum, and Cadmium in bee honey, in turn, is the reason for the increase in their content in the tissues of honey bees. In particular, it was established that in the tissues of the abdomen (respectively 163.76 g·10⁻³/kg of raw mass and 191.91 against 127.61), breast (80.10 and 97.24 against 65.27), and head (respectively 100.32 and 119.32 against 81.24 g·10⁻³/kg of raw weight) of honey bees of the foothill and forest-steppe

zones, compared to the mountain zone, there is a higher total content of the studied heavy metals (Tables 3, 4 and 5). The level of dangerous elements of the first class of toxicity - Lead (by 1.33-4.00 times) and Cadmium (by 1.78-4.00 times) - is especially significant in the tissues of honey bees of the foothills and forest-steppe zones, compared to the conditionally clean mountain environment). The concentration of the element of the second toxicity class - Chromium - also increases noticeably (by 1.18-1.60 times).

Table 3. The content of heavy metals, including toxic ones, in the abdominal tissues of honey bees in different natural zones of the Carpathian region, g·10⁻³/kg raw weight (M ± m, n = 3)

Metal and its symbol	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Ferrum, Fe	46.48 ± 1.046	63.72 ± 1.220***	77.03 ± 1.630***
Zinc, Zn	77.08 ± 1.190	91.32 ± 1.536**	104.24 ± 2.060***
Cuprum, Cu	0.34 ± 0.012	0.47 ± 0.014**	0.59 ± 0.014***
Cobalt, Co	0.31 ± 0.009	0.36 ± 0.014*	0.43 ± 0.017**
Chromium, Cr	2.43 ± 0.070	3.12 ± 0.082*	3.78 ± 0.112**
Nickel, Ni	2.43 ± 0.035	3.40 ± 0.067***	4.13 ± 0.059***
Lead, Pb	0.88 ± 0.035	1.21 ± 0.038**	1.50 ± 0.046***
Cadmium, Cd	0.09 ± 0.009	0.16 ± 0.006**	0.21 ± 0.012**

Table 4. The content of heavy metals, including toxic ones, in the breast tissues of honey bees in different natural zones of the Carpathian region, g·10⁻³/kg of raw mass (M ± m, n = 3)

Metal and its symbol	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Ferrum, Fe	36.10 ± 0.931	42.18 ± 0.812**	49.06 ± 0.555***
Zinc, Zn	17.51 ± 0.587	23.50 ± 0.625**	31.46 ± 0.507***
Cuprum, Cu	1.93 ± 0.041	2.82 ± 0.061***	3.17 ± 0.070***
Cobalt, Co	1.84 ± 0.035	2.19 ± 0.077*	2.40 ± 0.049***
Chromium, Cr	3.08 ± 0.046	3.51 ± 0.058**	4.34 ± 0.186**
Nickel, Ni	4.00 ± 0.049	4.80 ± 0.085**	5.47 ± 0.128***
Lead, Pb	0.78 ± 0.023	1.04 ± 0.068*	1.25 ± 0.038***
Cadmium, Cd	0.03 ± 0.003	0.06 ± 0.003*	0.09 ± 0.003**

Table 5. The content of heavy metals, including toxic ones, in the tissues of the head of honey bees in different natural zones of the Carpathian region, g·10⁻³/kg of raw mass (M ± m, n = 3)

Metal and its symbol	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Ferrum, Fe	26.79 ± 0.607	31.16 ± 0.979*	36.59 ± 1.324**
Zinc, Zn	30.13 ± 0.630	36.30 ± 0.564**	44.00 ± 0.280***
Cuprum, Cu	8.39 ± 0.319	14.35 ± 0.417***	17.66 ± 0.400***
Cobalt, Co	7.22 ± 0.055	7.56 ± 0.078*	7.97 ± 0.084**
Chromium, Cr	6.87 ± 0.098	8.15 ± 0.117**	9.47 ± 0.254***
Nickel, Ni	1.00 ± 0.055	1.47 ± 0.041**	1.78 ± 0.058**
Lead, Pb	0.81 ± 0.030	1.25 ± 0.044**	1.73 ± 0.050***
Cadmium, Cd	0.03 ± 0.003	0.08 ± 0.003***	0.12 ± 0.006***

It should also be noted that the tissues of the abdomen of honeybees are much more active accumulators of heavy metals than the tissues of the

chest and head. Practically all heavy metals are accumulated in 1.61-2.04 times greater quantities in

the tissues of the abdomen of honey bees than in the tissues of the chest and head.

Heavy metals at a physiologically determined level are involved in metabolic processes and the content of fatty acids in the body tissues of bees. In particular, cobalt-initiated protein synthesis in bee tissues, due to the activation of transport and information nucleic acids (Osman et al., 2021), is usually accompanied by the accumulation of polyunsaturated fatty acids necessary for the construction of cytoplasmic and cell membranes.

Since it is part of 9-desaturase, cuprum in the tissues of the body of bees at a physiologically determined level contributes to the formation of palmitic and stearic monounsaturated fatty acids of the omega-7 (palmitoleic) and omega-9 (oleic) families, respectively (Di et al., 2020; Takic et al., 2021).

Linoleic and linolenic acids, sequentially synthesized in plant tissues from oleic acid, are considered indispensable for bees and, therefore, must enter their bodies with food (Hajiahmadi et al., 2020; Hsu et al., 2021; Takic et al., 2021). In the tissues of bees from linoleic and linolenic acids because zinc is part of 2-, 3-, 4-, 5- and 6-desaturases, even longer-chain and more unsaturated fatty acids, respectively, of the omega-6 families (eicosatriene, eicosatetraenoic-arachidonic, docosadiene, and docosatetraenoic) and omega-3 (eicosapentaenoic, docosatrienoic, docosapentaenoic, and docosahexaenoic) (Takic et al., 2021; Hsu et al., 2021).

Longer chains and more unsaturated fatty acids of the omega-3 and omega-6 families are precious for the bee body. As mentioned above, polyunsaturated fatty acids in the body tissues of bees are a source for constructing cytoplasmic and cell membranes and synthesizing biologically active derivatives - prostaglandins, thromboxanes, and leukotrienes (Trinkl et al., 2020; Mărgăoan et al., 2021; Matuszewska et al., 2021). Thus, polyunsaturated fatty acids of the omega-3 and omega-6 families affect the health and vitality of bees.

Precious polyunsaturated fatty acids for bees are fatty acids of the omega-3 family (eicosapentaenoic, docosatrienoic, docosapentaenoic, and docosahexaenoic) (Trinkl et al., 2020; Matuszewska et al., 2021; Takic et al., 2021). These acids in the body tissues of bees are the initiators of synthesizing very strong and fast-acting anti-inflammatory peptide substances - cytokines (Ranneh et al., 2021).

Based on the above, our next task was to study the concentration of esterified fatty acids in the tissues of the abdomen, breast, and head of honey bees obtained from hives located in the mountain, foothill, and forest-steppe zones of the Carpathian region.

The content of esterified forms of saturated fatty acids with an even (caprylic, capric, lauric, myristic, palmitic, and stearic) and odd (pentadecanic) amount of carboxylic acids was studied in the tissues of the abdomen, breast, and head of honey bees of the foothills and forest-steppe zones of the Carpathian region, compared to the mountain zone. Atoms in the chain, monounsaturated fatty acids of the omega-7 (palmitoleic) and omega-9 (oleic and eicosanoic) families, and polyunsaturated fatty acids of the omega-3 (linolenic, eicosapentaenoic, docosatrienoic, docosapentaenoic and docosahexaenoic) and omega-6 (linoleic, eicosadienoic) families, eicosatrienoic, eicosatetraenoic-arachidonic, docosadiene and docosatetraenoic). Data on their content are presented in Tables 6, 7, and 8.

It should be noted that the most significant amount of esterified forms of fatty acids is found in the tissues of the abdomen, breast, and head of honey bees (Desbois & Smith, 2010; Giri et al., 2018; Burdge, 2018; Saranchuk, 2020; Corby-Harris et al., 2021). Esterified forms of fatty acids in the mentioned tissues of honey bees are included in the composition of phospholipids, esterified cholesterol, mono-, di- and triacylglycerols (Desbois & Smith, 2010; Corby-Harris et al., 2021). It was established (Tables 6, 7 and 8) that the total concentration of esterified forms of fatty acids in the tissues of the abdomen, chest and head of honey bees obtained from hives located in the foothills (respectively 17.67, 18.57 and 18.33 g/kg of raw mass) and forest-steppe (16.98, 17.69 and 17.31) zones of the Carpathian region, compared with the tissues of the abdomen, breast and head of honey bees selected from hives located in the mountain zone (21.72, 22, 69 and 23.17), due to the lower level of saturated fatty acids in their composition (2.24, 2.32 and 1.99 and 2.09, 2.12 and 1.96 versus 2.93, 2.97 and 2.97) and an odd (0.07, 0.07 and 0.09 and 0.06, 0.05 and 0.08 versus 0.09, 0.09 and 0.13) number of carbon atoms in the chain, of monounsaturated fatty acids of the omega-7 (0.05, 0.05 and 0.05 and 0.04, 0.04 and 0.04 versus 0.08, 0.07 and 0.07) and omega-9 (3.53, 3.31 and 3.12 and 3.36, 3.20 and 3.03 vs. 4.12, 4.00 and 3.78) and polyunsaturated fatty acids of omega-3 families (6.18, 6, 62 and 7.29 and 5.97, 6.42 and 6.98 against 7.61, 8.01 and 9.10) and about mega-6 (respectively 5.60, 6.10 and 5.79 and 5.46, 5.86 and 5.22 against 6.89, 7.55 and 7.12 g/kg raw weight).

In terms of energy, lipids are much more valuable than proteins and carbohydrates (Ruedenauer et al., 2021; Stabler et al., 2021). Literature sources indicate that the greater the number of fatty acids contained in tissues, the

greater their energy value for the body of honey bees (Ruedenauer et al., 2021). It was established (Tables 6, 7, and 8) that the most significant amount of esterified forms of saturated, monounsaturated, and polyunsaturated fatty acids are contained in the tissues of the abdomen, breast, and head of honey bees in the mountainous zone of the

Carpathian region (respectively 21.72, 22.69 and 23, 17 g/kg of raw weight), a smaller amount of them is in the tissues of the abdomen, chest, and head of honey bees in the foothills (17.67, 18.57 and 18.33), and even less in the forest-steppe (respectively 16.98, 17, 69 and 17.31 g/kg raw weight).

Table 6. The content of esterified fatty acids in the abdominal tissues of honey bees in different natural zones of the Carpathian region at the beginning of the summer period, g/kg of raw mass ($M \pm m$, $n = 3$)

Acid and its code	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Caprylic acid, 8:0	0.08 ± 0.003	0.04 ± 0.003**	0.03 ± 0.003**
Capric acid, 10:0	0.05 ± 0.003	0.03 ± 0.003*	0.02 ± 0.003**
Lauric acid, 12:0	0.06 ± 0.003	0.04 ± 0.003*	0.03 ± 0.003**
Myristic acid, 14:0	0.07 ± 0.003	0.05 ± 0.003*	0.04 ± 0.003**
Pentadecanoic acid, 15:0	0.09 ± 0.003	0.07 ± 0.003*	0.06 ± 0.003**
Palmitic acid, 16:0	1.20 ± 0.028	0.93 ± 0.024**	0.92 ± 0.012***
Palmitoleic acid, 16:1	0.08 ± 0.003	0.05 ± 0.003*	0.04 ± 0.003**
Stearic acid, 18:0	1.30 ± 0.032	1.04 ± 0.045**	0.95 ± 0.024***
Oleic acid, 18:1	3.91 ± 0.085	3.36 ± 0.031**	3.21 ± 0.059**
Linoleic acid, 18:2	2.79 ± 0.074	2.28 ± 0.080**	2.23 ± 0.038**
Linolenic acid, 18:3	4.05 ± 0.113	3.31 ± 0.052**	3.20 ± 0.067**
Arachinic acid, 20:0	0.17 ± 0.007	0.11 ± 0.005**	0.10 ± 0.006**
Eicosaenoic acid, 20:1	0.21 ± 0.006	0.17 ± 0.003*	0.15 ± 0.003**
Eicosadienoic acid, 20:2	0.21 ± 0.009	0.17 ± 0.003*	0.15 ± 0.003**
Eicosatrienoic acid, 20:3	0.31 ± 0.009	0.23 ± 0.009**	0.21 ± 0.006**
Arachidonic acid, 20:4	3.18 ± 0.076	2.67 ± 0.035**	2.60 ± 0.032**
Eicosapentaenoic acid, 20:5	2.17 ± 0.071	1.74 ± 0.023**	1.71 ± 0.017**
Docosadienoic acid, 22:2	0.28 ± 0.012	0.21 ± 0.006**	0.19 ± 0.007**
Docosatrienoic acid, 22:3	0.32 ± 0.010	0.23 ± 0.006**	0.21 ± 0.006**
Docosatetraenoic acid, 22:4	0.32 ± 0.009	0.25 ± 0.006**	0.23 ± 0.007**
Docosapentaenoic acid, 22:5	0.53 ± 0.015	0.43 ± 0.006**	0.40 ± 0.007**
Docosahexaenoic acid, 22:6	0.54 ± 0.009	0.47 ± 0.015*	0.45 ± 0.007**

Table 7. The content of esterified fatty acids in the breast tissues of honey bees in different natural zones of the Carpathian region at the beginning of the summer period, g/kg of raw weight ($M \pm m$, $n = 3$)

Acid and its code	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Caprylic acid, 8:0	0.03 ± 0.003	0.02 ± 0.003*	0.01 ± 0.000**
Capric acid, 10:0	0.04 ± 0.003	0.02 ± 0.003*	0.01 ± 0.003**
Lauric acid, 12:0	0.05 ± 0.003	0.03 ± 0.003*	0.02 ± 0.006*
Myristic acid, 14:0	0.09 ± 0.003	0.06 ± 0.003**	0.04 ± 0.003***
Pentadecanoic acid, 15:0	0.09 ± 0.006	0.07 ± 0.003*	0.05 ± 0.003**
Palmitic acid, 16:0	1.29 ± 0.026	1.08 ± 0.022**	0.97 ± 0.098**
Palmitoleic acid, 16:1	0.07 ± 0.003	0.05 ± 0.003*	0.04 ± 0.003**
Stearic acid, 18:0	1.30 ± 0.032	1.02 ± 0.050**	0.97 ± 0.047**
Oleic acid, 18:1	3.74 ± 0.066	3.13 ± 0.073**	3.04 ± 0.084**
Linoleic acid, 18:2	2.95 ± 0.064	2.37 ± 0.049**	2.28 ± 0.052**
Linolenic acid, 18:3	3.58 ± 0.118	3.02 ± 0.058*	2.92 ± 0.04**
Arachinic acid, 20:0	0.17 ± 0.006	0.11 ± 0.006**	0.10 ± 0.006**
Eicosaenoic acid, 20:1	0.26 ± 0.014	0.18 ± 0.009**	0.16 ± 0.012**
Eicosadienoic acid, 20:2	0.28 ± 0.009	0.20 ± 0.015*	0.19 ± 0.015*
Eicosatrienoic acid, 20:3	0.21 ± 0.009	0.16 ± 0.006**	0.15 ± 0.006**
Arachidonic acid, 20:4	3.51 ± 0.082	3.00 ± 0.050**	2.80 ± 0.128**
Eicosapentaenoic acid, 20:5	2.43 ± 0.050	1.98 ± 0.041**	1.97 ± 0.044**
Docosadienoic acid, 22:2	0.23 ± 0.009	0.18 ± 0.007*	0.17 ± 0.007**

Docosatrienoic acid, 22:3	0.27 ± 0.012	0.19 ± 0.007**	0.17 ± 0.009**
Docosatetraenoic acid, 22:4	0.37 ± 0.014	0.29 ± 0.009**	0.27 ± 0.009**
Docosapentaenoic acid, 22:5	0.81 ± 0.034	0.67 ± 0.009*	0.63 ± 0.019*
Docosahexaenoic acid, 22:6	0.92 ± 0.029	0.76 ± 0.019**	0.73 ± 0.017**

Table 8. The content of esterified fatty acids in the tissues of the head of honey bees in different natural zones of the Carpathian region at the beginning of the summer period, g/kg of raw weight ($M \pm m$, $n = 3$)

Acid and its code	Natural zones of the Carpathian region		
	Mountain	Foothills	Forest-steppe
Caprylic acid, 8:0	0.03 ± 0.003	0.02 ± 0.003*	0.01 ± 0.003*
Capric acid, 10:0	0.04 ± 0.003	0.02 ± 0.003*	0.01 ± 0.003**
Lauric acid, 12:0	0.05 ± 0.003	0.03 ± 0.003*	0.02 ± 0.003**
Myristic acid, 14:0	0.09 ± 0.003	0.06 ± 0.003**	0.05 ± 0.003**
Pentadecanoic acid, 15:0	0.13 ± 0.003	0.09 ± 0.003**	0.08 ± 0.007**
Palmitic acid, 16:0	1.52 ± 0.084	0.98 ± 0.030*	0.94 ± 0.029**
Palmitoleic acid, 16:1	0.07 ± 0.003	0.05 ± 0.003*	0.04 ± 0.003**
Stearic acid, 18:0	1.05 ± 0.038	0.86 ± 0.014*	0.83 ± 0.015**
Oleic acid, 18:1	3.54 ± 0.089	2.95 ± 0.045**	2.87 ± 0.026**
Linoleic acid, 18:2	2.77 ± 0.068	2.15 ± 0.036**	2.03 ± 0.037***
Linolenic acid, 18:3	3.54 ± 0.075	3.00 ± 0.054**	2.92 ± 0.046**
Arachidonic acid, 20:0	0.18 ± 0.012	0.11 ± 0.007**	0.10 ± 0.007**
Eicosaenoic acid, 20:1	0.24 ± 0.012	0.17 ± 0.006**	0.16 ± 0.003**
Eicosadienoic acid, 20:2	0.19 ± 0.006	0.14 ± 0.006**	0.13 ± 0.006**
Eicosatrienoic acid, 20:3	0.13 ± 0.003	0.10 ± 0.003**	0.09 ± 0.003**
Arachidonic acid, 20:4	3.35 ± 0.072	2.85 ± 0.087*	2.44 ± 0.042***
Eicosapentaenoic acid, 20:5	2.47 ± 0.081	1.94 ± 0.058**	1.87 ± 0.037**
Docosadienoic acid, 22:2	0.22 ± 0.009	0.17 ± 0.006**	0.16 ± 0.006**
Docosatrienoic acid, 22:3	0.23 ± 0.006	0.18 ± 0.006**	0.16 ± 0.006**
Docosatetraenoic acid, 22:4	0.46 ± 0.014	0.38 ± 0.006**	0.37 ± 0.006**
Docosapentaenoic acid, 22:5	1.24 ± 0.038	0.91 ± 0.029**	0.86 ± 0.032**
Docosahexaenoic acid, 22:6	1.62 ± 0.058	1.26 ± 0.029**	1.17 ± 0.038**

Esterified forms of polyunsaturated fatty acids of the omega-3 and omega-6 families in the composition of phospholipids are included in the structure of cellular and cytoplasmic membranes of the body and ensure their functional activity and, ultimately, the vital activity of bees (Trinkl et al., 2020). At the same time, the esterified form of linoleic acid and its longer-chain and unsaturated fatty acids in the body of bees is the initiator of pro-inflammatory processes (Mărgăoan et al., 2021), and linoleic acid and its longer-chain and unsaturated fatty acids are anti-inflammatory (Ranneh et al., 2021). Esterified forms of linoleic and linolenic acids and their longer-chain and unsaturated fatty acids act on the bees' bodies through the corresponding pro-inflammatory and anti-inflammatory peptide cytokines (Ranneh et al., 2021).

It was recorded (Tables 6, 7, and 8) that the most significant amount of esterified forms of polyunsaturated fatty acids of the omega-3 and omega-6 families are contained in the tissues of the abdomen, breast, and head of honey bees of

the mountain zone of the Carpathian region (respectively 7.61, 8.01 and 9.10 and 6.89, 7.55 and 7.55 g/kg of raw weight), a smaller number of them are in the tissues of the abdomen, chest, and head of honey bees in the foothill zone (6.18, 6.62 and 7, 29 and 5.60, 6.10 and 5.79), even less in the forest-steppe (respectively 5.97, 6.42 and 6.98 and 5.46, 5.86 and 5.22 g/kg raw weight). The mentioned esterified forms of polyunsaturated fatty acids of the omega-6 and omega-3 families in the exact quantities in bee tissues are initiators of pro-inflammatory and anti-inflammatory processes, respectively.

Esterified forms of polyunsaturated fatty acids linolenic and linoleic in the tissues of the bee body are primarily the precursors of a whole series of longer-chain and more unsaturated fatty acids, respectively, of the families of omega-3 (eicosapentaenoic, docosatrienoic, docosapentaenoic and docosahexaenoic) and omega-6 (eicosatrienoic, eicosatetraenoic-arachidonic, docosadiene and docosaterpene). Zinc takes an active part in the reactions of

conversion of linoleic and linolenic acids into their longer-chain and more unsaturated derivatives in bee tissues, as it is part of 2-, 3-, 4-, 5-, and 6-desaturases (Di et al., 2020; Hsu et al., 2021). At the same time, the ratio of the content of esterified forms of linolenic and linoleic acids to the content of their longer-chain and unsaturated fatty acids of the omega-3 and omega-6 families, respectively, indicates the intensity of transformation of the less functionally and biologically active former into more active latter (Hsu et al., 2021).

It was established (Tables 6, 7, and 8) that the esterified form of linolenic and linoleic acids in the abdominal tissues of honey bees kept in the foothills and forest-steppe zones of the Carpathian region, compared to the mountain ones, are less wholly transformed into their longer-chain and unsaturated fatty acids of the families omega-3 (1.15 and 1.16 vs. 1.14, respectively) and omega-6 (0.69 and 0.69 vs. 0.68, respectively). The same direction of transformations of the esterified form of linolenic acid is observed in the breast tissues of honey bees (0.82 and 0.83 versus 0.81). At the same time, in the breast tissues of honey bees of the forest-steppe zone of the Carpathian region, compared to breast tissues of honey bees of the mountain zone, the intensity of conversion of the esterified form of linoleic acid into its longer-chain and unsaturated fatty acids of the omega-6 family is sharply reduced (0.64 vs. 0.60). A similar direction of transformations of the esterified form of linoleic acid is observed in the tissues of the head of honey bees of the foothill zone of the Carpathian region (0.64 versus 0.59). At the same time, in the tissues of the head of honey bees, which are kept in the foothills and especially in the forest-steppe zones of the Carpathian region, compared to the mountains, the intensity of the conversion of the esterified form of linolenic acid into its longer-chain and unsaturated fatty acids of the omega-3 family (respectively 0.70 and 0.72 vs. 0.64).

The above-mentioned indicates the negative influence of several heavy metals present, primarily toxic Lead and Cadmium, on the synthesis in the body tissues of honeybees of very active biologically and functionally longer chain and unsaturated fatty acids of the omega-3 and omega-6 families.

The low total content of esterified polyunsaturated and monounsaturated fatty acids of the omega-3, omega-6, omega-7, and omega-9 families can contribute to the increase in the fragility of bee comb walls through the bee body, in particular the wax glands (Didaras et al., 2020; Castaños et al., 2022). At the same time, the shallow content of the above fatty acids in the tissues of the abdomen, chest, and head of honeybees can cause a decrease in the permeability of its structural components to water and water-soluble substances and thereby inhibit the intensity of metabolic processes (Bakour et al., 2022; Castaños et al., 2022). It can also contribute to a decrease in the functional activity of cellular and cytoplasmic membranes of the body of honey bees and thereby inhibit their vital activity (Corby-Harris et al., 2021; Ricigliano et al., 2021; Wang et al., 2021).

It was established that the total content of esterified polyunsaturated and monounsaturated fatty acids of the omega-3, omega-6, omega-7, and omega-9 families in the tissues of the abdomen, breast, and head of honey bees obtained from hives located in the foothills (respectively 15.36, 16.08 and 16.25 g/kg of raw weight) and forest-steppe (14.83, 15.52 and 15.27) zones of the Carpathian region, compared to the tissues of the abdomen, breast, and head of honey bees selected from hives located in the mountain zone (18.70, 19.63 and 20.07 g/kg raw weight, respectively), is smaller (Tables 6, 7 and 8).

Saturated, monounsaturated, and polyunsaturated fatty acids caprylic, capric, lauric, myristic, palmitoleic, oleic, linoleic, linolenic, eicosaenic, eicosadiene, eicosatriene, eicosatetraenoic-arachidonic, eicosapentaenoic, docosadiene, docosatrienoic, docosatetraenoic, docosapentaenoic, and docosahexaenoic provide antibacterial protection of the body honey bees (Didaras et al., 2020; Rothman et al., 2020; Mărgăoan et al., 2021; Bakour et al., 2022). In particular, the high antibacterial activity of the above acids against bee rot was found (Rothman et al., 2020).

Caprylic and, to a lesser extent, capric and, to a lesser extent, lauric and, to a lesser extent, myristic acids have an antimicrobial effect due to their high ability to reduce the concentration of hydrogen ions (Didaras et al., 2020;

Rothman et al., 2020; Bakour et al., 2022), and palmitooleic, oleic, linoleic, linolenic, eicosaic, eicosadiene, eicosatriene, eicosatetraeno-arachidone, eicosapentaeno, docosadiene, docosatrieno, docosatetraenoic, docosapentaenoic, and docosahexaenoic - with the growth of this series, increase the surface activity of the tissues of microorganisms and thereby strongly suppress their vital activity under the normal osmotic pressure of the surrounding environment (Wang et al., 2021; Mărgăoan et al., 2021).

It was established that the total content of esterified caprylic, capric, lauric, myristic, palmitoleic, oleic, linoleic, linolenic, eicosenoic, eicosadienoic, eicosatrienoic, eicosatetraenoic-arachidonic, eicosapentaenoic, docosadienoic, docosatrienoic, docosatetraenoic, docosapentaenoic, and docosahexaenoic acids, which provide antibacterial protection of the bee organism, in the tissues of the abdomen, chest, and head of honey bees obtained from hives located in the foothills (respectively 15.36, 16.31 and 16.38 g/kg of raw mass) and forest-steppe (15.31, 15.60 and 15.36) zones of the Carpathian region, compared to the tissues of the abdomen, chest, and head of honey bees selected from hives located in the mountain zone (respectively 19.16, 19.84 and 20.28 g/kg of raw mass), there is less (Tables 5, 6 and 7). This significantly affects, as evidenced by the scientific literature (Didaras et al., 2020; Mărgăoan et al., 2021; Wang et al., 2021), the antibacterial and antifungal activity of the body tissues of honey bees.

Thus, as a result of an increase in the artificial load on the environment and the accumulation of heavy metals, primarily toxic, in the components of the ecosystem, the energy, structural, biological, and antimicrobial value of the central mass of long-chain fatty acids for the body of bees decreases.

The decrease in the content of esterified forms of fatty acids in the tissues of honey bees kept in hives located in the foothills and especially the forest-steppe zones of the Carpathian region, compared to the tissues of honey bees selected from hives located in the mountainous zone, is associated with their transition to anionic form. This is due primarily to the

binding of fatty acids by heavy metal cations (Giri et al., 2018).

It was established that the honey productivity of worker bees in the mountainous zone of the Carpathian region is at the level of 46.6 ± 0.95 kg, in the foothills - 36.6 ± 1.04 , $p < 0.05$, and in the forest-steppe - 31.2 ± 0.56 kg, $p < 0.01$ per bee colony per season. The honey productivity of worker bees decreases due to the high level of heavy metals, including toxic but low - esterified fatty acids, in the tissues. Other scientists also point to the negative influence of territories polluted with heavy metals on the productive characteristics of honey bees (Gizaw et al., 2020; Monchanin et al., 2021).

All over the world, there is a search for means of bioindication of the ecological state of the environment (Komarova, 2018; Costa et al., 2021). This is because heavy metals, like other environmental pollutants, have different transfer rates from the soil to the root system, from the root system to the stem, from the stem to the inflorescence, and from the inflorescence to pollen and nectar.

It was previously indicated (Saranchuk & Ravis, 2008) that *Taraxacum officinale* Wigg pollen can serve as a bioindicator of the ecological state of the environment in the conditions of the Carpathian region due to the optimal content of heavy metals and fatty acids. The positive thing about this bioindicator is that it allows to determine different levels of accumulation of heavy metals and fatty acids and thus gives more information. The body tissues of honey bees can also serve as a bioindicator of the ecological state of the environment in terms of the content of heavy metals and esterified forms of fatty acids.

CONCLUSIONS

The total content of the studied heavy metals in the tissues of honey bees in the foothills and forest-steppe zones of the Carpathian region, compared to the mountain zone, is 1.23-1.50 times higher. The level of dangerous elements of the first class of toxicity - Lead (by 1.33-4.00 times) and Cadmium (by 1.78-4.00 times) - is especially significant in the tissues of honey bees of the foothills and forest-steppe zones, compared to the conditionally clean mountain environment).

In the Carpathian region, all heavy metals accumulate 1.61-2.04 times more in the abdominal

tissues of honey bees than in the breast and head tissues.

In the direction from the mountain to the foothills and further to the forest-steppe zone of the Carpathian region, a decrease in the content of esterified forms of fatty acids is observed in the tissues of honey bees (21.72-23.17, 17.67-18.57 and 16.98-17.69 g/kg raw mass).

Esterified forms of linolenic and linoleic acids in the abdominal tissues of honey bees kept in the foothills and forest-steppe zones of the Carpathian region, compared to the mountain ones, are less completely converted into their longer-chain and unsaturated fatty acids of the omega-3 families (respectively 1, 15 and 1.16 vs. 1.14) and omega-6 (0.69 and 0.69 vs. 0.68, respectively). At the same time, the intensity of conversion of the esterified form of linoleic acid into its longer chain and unsaturated derivatives of the omega-6. Also, the intensity of the conversion of the esterified form of linolenic acid into its longer-chain and unsaturated fatty acids of the omega-3 family in the tissues of the head of honey bees in the foothills and especially in the forest-steppe zones (0.70 and 0.72 vs. 0.64, respectively) is sharply reduced.

As a result of the accumulation of heavy metals in tissues, energy (by 21.0-44.6%), structural (21.2-45.7), biological (8.7-24.7), and antimicrobial (by 21.6-32.1%) values of esterified forms of fatty acids for the body of honey bees kept in hives located in the foothills and especially forest-steppe zones of the Carpathian region, compared to the tissues of bees kept in apiaries located in the mountainous zone.

The honey productivity of worker bees per beehive per season is lower in the foothills and forest-steppe zones of the Carpathian region, compared to the mountain, by 27.3% and 49.4%, respectively.

Body tissues of honey bees can serve as a bioindicator of the ecological state of the environment in terms of the content of heavy metals and esterified forms of fatty acids

REFERENCES

- Al-Kahtani, S. N., Taha, E. K. A., Farag, S. A., Taha, R. A., Abdou, E. A., & Mahfouz, H. M. (2021). Harvest Season Significantly Influences the Fatty Acid Composition of Bee Pollen. *Biology*, 10(6), 495–504.
- Bakour, M., Laaroussi, H., Ousaaid, D., El Ghouizi, A., Es-Safi, I., Mechchate, H., & Lyoussi, B. (2022). Bee Bread as a Promising Source of Bioactive Molecules and Functional Properties: An Up-To-Date Review. *Antibiotics*, 11(2), 1–39.
- Bashchenko, M. I., Boiko, O. V., Honchar, O. F., Gutyj, B. V., Lesyk, Y. V., Ostapyuk, A. Y., Kovalchuk, I. I., & Leskiv, Kh. Ya. (2020). The effect of milk thistle, metiphen, and silimevit on the protein-synthesizing function of the liver of laying hens in experimental chronic cadmium toxicosis. *Ukrainian Journal of Ecology*, 10(6), 164–168.
- Briffa, J., Sinagra, E., & Blundell, R. (2020). Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon*, 6(9), 1–26.
- Burdge, G. C. (2018). *Polyunsaturated fatty acid metabolism*. Cambridge, USA: Academic Press et AOCS Press.
- Butsiak, H. A., Butsiak, V. I., Gutyj, B. V., Kalyn, B. M., Muzyka, L. I., Stadnytska, O. I., Luchyn, I. S., Rozputnii, O. I., Kachan, L. M., Melnichenko, Yu. O., Sliusarenko, S. V., Bilkevich, V. V., & Leskiv, K. Y. (2021). Migration of mobile forms of heavy metals into the vegetative mass of plants under local human-caused load. *Ukrainian Journal of Ecology*, 11 (1), 239–343.
- Castaños, C., Boyce, M., Millar, H., Lawler, N., Bates, T., & Grass, J. (2022). Mobilization of Lipids Underpins Honey Bee and Colony Health During Limited Supplementary Feeding. *Research Square*, 1–28.
- Corby-Harris, V., Bennett, M. M., Deeter, M. E., Snyder, L., Meador, C., Welchert, A. C., Hoffman, A., Obernesser, B. T., & Carroll, M. J. (2021). Fatty acid homeostasis in honey bees (*Apis mellifera*) fed commercial diet supplements. *Apidologie*, 52, 1195–1209.
- Costa, A., Veca, M., Barberis, M., Ciceronegri, L., & Tangorra, F. M. (2021). Predicting atmospheric cadmium and lead using honeybees as atmospheric heavy metals pollution indicators. Results of a monitoring survey in Northern Italy. *Italian Journal of Animal Science*, 20(1), 850–858.
- Desbois, A. P., & Smith, V. J. (2010). Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential. *Applied Microbiology and Biotechnology*, 85(6), 1629–1642.
- Di, N., Zhang, K., Hladun, K. R., Rust, M., Chen, Y.-F., Zhu, Z.-Y., Liu, T.-X., & Trumble, J. T. (2020). Joint effects of cadmium and copper on *Apis mellifera* forgers and larvae. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 237, 1–10.
- Didaras, N. A., Karatasou, K., Dimitriou, T. G., Amoutzias, G. D., & Mossialos, D. (2020). Antimicrobial Activity of Bee-Collected Pollen and Beebread: State of the Art and Future Perspectives. *Antibiotics*, 9(11), 811–840.
- El-Seedi, H. R., Eid, N., Abd El-Wahed, A. A., Rateb, M. E., Afifi, H. S., Algethami, A. F., Zhao, C., Al Naggari, Y., Alsharif, S. M., Tahir, H. E., Xu, B., Wang, K., & Khalifa, S. A. (2022). M. Honey Bee Products: Preclinical and Clinical Studies of Their Anti-inflammatory and Immunomodulatory Properties. *Frontiers in Nutrition*, 8, 761267. doi: 10.3389/fnut.2021.761267
- Giri, S., Rule, D. C., & Dillon, M. E. (2018). Fatty acid composition in native bees: Associations with thermal and feeding ecology. *Comparative Biochemistry and Physiology Part A*, 218, 70–79.
- Gizaw, G., Kim, Y. H., Moon, K. H., Choi, J. B., Kim, Y. H., & Park, J. K. (2020). Effect of environmental heavy metals on the expression of detoxification-

- related genes in honey bee *Apis mellifera*. *Apidologie*, 51, 664–674.
- Gutyj, B., Ostapiuk, A., Kachmar, N., Stadnytska, O., Sobolev, O., Binkevych, V., Petryshak, R., Petryshak, O., Kulyaba, O., Naumyuk, A., Nedashkivsky, V., Nedashkivska, N., Magrelo, N., Golodyuk, I., Nazaruk, N., & Binkevych, O. (2019). The effect of cadmium loading on protein synthesis function and functional state of laying hens' liver. *Ukrainian Journal of Ecology*, 9(3), 222–226.
- Gutyj, B. V., Ostapyuk, A. Y., Sobolev, O. I., Vishchur, V. J., Gubash, O. P., Kurtyak, B. M., Kovalskyi, Y. V., Darmohray, L. M., Hunchak, A. V., Tsisaryk, O. Y., Sheherbatyy, A. R., Farionik, T. V., Savchuk, L. B., Palyadichuk, O. R., Hrymak, K. (2019). Cadmium burden impact on morphological and biochemical blood indicators of poultry. *Ukrainian Journal of Ecology*, 9(1), 236–239.
- Gutyj, B., Stybel, V., Darmohray, L., Lavryshyn, Y., Turko, I., Hachak, Y., Sheherbatyy, A., Bushueva, I., Parchenko, V., Kaplaushenko, A., Krushelnyska, O. (2017). Prooxidant-antioxidant balance in the organism of bulls (young cattle) after using cadmium load. *Ukrainian Journal of Ecology*, 7(4), 589–596.
- Hajiahmadi, Z., Abedi, A., Wei, H., Sun, W., Ruan, H., Zhuge, Q., & Movahedi, A. (2020). Identification, evolution, expression, and docking studies of fatty acid desaturase genes in wheat (*Triticum aestivum* L.). *BMC Genomics*, 21(778), 1–20.
- Hsu, P.S., Wu, T.-H., Huang, M.Y., Wang, D.Y., & Wu, M.C. (2021). Nutritive Value of 11 Bee Pollen Samples from Major Floral Sources in Taiwan. *Foods*, 10(9), 2229–2244.
- Ibatullin, I. I., & Zhukorskyi, O. M. (2017). Methodology and organization of scientific research in animal husbandry: a guide. Kyiv, UA: Agrarian Science Publishing House.
- Klym, O., & Stadnytska, O. (2019). Heavy metals in the dandelion and apple tree pollen from the different terrestrial ecosystems of the Carpathian region. *Acta Scientiarum Polonorum Zootechnica*, 18(3), 15–20.
- Kobys, A. I., Chechet, O. M., Shuliak, S. V., Omelchun, Yu. A., Miahka, K. S., Marchenko, T. V., & Liniichuk, N. V. (2021). The problem of the spread of toxicants in animal husbandry and the environment. *Bulletin of the Sumy National Agrarian University. "Veterinary Medicine" series*, 3(54), 17–25.
- Komarova, I. (2018). *Taraxacum officinale* as bioindicator of heavy metal accumulation in soil. Danish Scientific Journal (DSJ). *Istedgade 1041650 København V Denmark*, 8, 10–12.
- Kovalskyi, Y. V., & Kyryliv, Y. I. (2011). Technology of obtaining beekeeping products. Lviv, UA: Lviv Polytechnic Publishing House.
- Kovalskyi, Y., Gucol, A., Gutyj, B., Sobolev, O., Kovalska, L., & Mironovych, A. (2018). Features of histology and histogenesis in the vital temperature range in the organism of honey bee (*Apis mellifera* L.) in the postembryonic period. *Ukrainian Journal of Ecology*, 8(2), 301–307.
- Kovka, N. O., & Nedashkivskyi, V. M. (2019). Duration and periods of flowering of the main nectarifera in conditions of the right-bank forest-steppe. *Livestock breeding of Ukraine*, 4, 36–39.
- Krainiukov, O., Kryvytska, I., & Cherkashyna, Yu. (2020). Assessment of the influence of heavy metals on the photosynthetic apparatus of plants. *Young scientist. Biological sciences*, 4(80), 244–252.
- Kryvyi, M. M., Zhukovets, O. I., & Dikhtiar, O. O. (2018). Assessment of honey-bearing resources of forest ecosystems based on their typology. *Agrarian science. Animal feeding and feed technology*, 2(101), 34–43.
- Lorelta, Y., Yong, R. N., & Thomas, H. R. (2015). Fate and Transport of Lead Pollution Along a Highway Corridor. *Geoenvironmental engineering*. doi: 10.1680/jgeimogacl.32774.0012.
- Mărgăoan, R., Özkök, A., Keskin, Ş., Mayda, N., Urcan, A. C., & Cornea-Cipcigan, M. (2021). Bee collected pollen as a value-added product rich in bioactive compounds and unsaturated fatty acids: A comparative study from Turkey and Romania. *LWT*, 149, 111925. doi: 10.1016/j.lwt.2021.111925
- Matin, G., Kargar, N., & Buyukisik, H. B. (2016). Biomonitoring of cadmium, lead, arsenic and mercury in industrial districts of Izmir, Turkey by using honey bees, propolis and pine tree leaves. *Ecological Engineering*, 90(5), 331–335.
- Matuszewska, E., Klupezyńska, A., Maciołek, K., Kokot, Z. J., & Matysiak, J. (2021). Multielemental Analysis of Bee Pollen, Propolis, and Royal Jelly Collected in West-Central Poland. *Molecules*, 26(9), 1–18.
- Monchanin, C., Drujont, E., Devaud, J.-M., Lihoreau, M., & Barron, A. B. (2021). Metal pollutants have additive negative effects on honey bee cognition. *Journal of Experimental Biology*, 224, 1–9.
- Osman, D., Cooke, A., Young, T. R., Deery, E., Robinson, N., & Warren, M. J. (2021). The requirement for cobalt in vitamin B₁₂: A paradigm for protein metalation. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1868(1), 118896. doi: 10.1016/j.bbamcr.2020.118896
- Prychepa, M., Hrynevych, N., Martseniuk, V., Potrokhov, O., Vodianskyi, O., Khomiak, O., Rud, O., Kytsokon, L., Sliusarenko, A., Dunaievskia, O., Gutyj, B., Pukalo, P., Honcharenko, V., Yevtukh, L., Bozhyk, L., Prus, V., & Makhorin, H. (2021). Rudd (*Scardinius erythrophthalmus* L., 1758) as a bioindicator of anthropogenic pollution in freshwater bodies. *Ukrainian Journal of Ecology*, 11(2), 253–260.
- Purać, J., Nikolić, T. V., Kojić, D., Čelić, A. S., Plavša, J. J., Blagojević, D. P., & Petri, E. T. (2019). Identification of a metallothionein gene in honey bee *Apis mellifera* and its expression profile in response to Cd, Cu and Pb exposure. *Molecular Ecology*, 28(4), 731–745.
- Ranneh, Y., Akim, A. M., Hamid, H. A., Khazaai, H., Fadel, A., Zakaria, Z. A., Albuja, M., Fadzelly, M., & Bakar, A. (2021). Honey and its nutritional and anti-inflammatory value. *BMC Complementary Medicine and Therapies*, 21(30), 1–17.
- Razanov, S. F., Didur, I. M., & Pervachuk, M. V. (2015). Effectiveness of reduction of soil pollution by lead and cadmium by bee pollination of agricultural crops

- in the conditions of their mineral nutrition. *Agriculture and forestry*, 2, 94–101.
- Ricigliano, V. A., Dong, C., Richardson, L. T., Donnarumma, F., Williams, S. T., Solouki, T., & Murray, K. K. (2021). Honey Bee Proteome Responses to Plant and Cyanobacteria (blue-green algae) Diets. *ACS Food Science & Technology*, 1(1), 17–26.
- Rivis, Y., Zaborski, D., Gutyj, B., Hopanenko, O., Diachenko, O., Stadnytska, O., Klum, O., Saranchuk, I., Bratyuk, V., & Fedak, V. (2022). Quantitative and simultaneous gas chromatographic determination of various forms long-chain fatty acids in biological material. *Scientific Papers. Series D. Animal Science*, 65(2), 24–29.
- Rothman, J. A., Russell, K. A., Leger, L., McFrederick, Q. S., & Graystock, P. (2020). The direct and indirect effects of environmental toxicants on the health of bumble bees and their microbiomes. *Proceedings of the Royal Society B: Biological Sciences*. doi: 10.1101/2020.04.24.060228
- Ruedenaucr, F. A., Biewer, N. W., Nebauer, C. A., Scheiner, M., Spaethe, J., & Leonhardt, S. D. (2021). Honey Bees Can Taste Amino and Fatty Acids in Pollen, but Not Sterols. *Frontiers in Ecology and Evolution*, 9, 684175.
- Saranchuk, I. I. (2020). Level of non-esterified fatty acids in abdominal tissues and productive traits of bees fed different amounts of linseed oil. *Foothill and mountain agriculture and animal husbandry*, 67(II), 253–264.
- Saranchuk, I. I., & Rivis, Y. F. (2008). Fatty acid composition of bee pollen depending on environmental conditions. *Biology of animals*, 10(1, 2), 236–244.
- Saranchuk, I. I., Vishchur, V. Ya., Gutyj, B. V., & Klim, O. Y. (2021). Effect of various amounts of sunflower oil in feed additives on breast tissues' functional condition, reproductivity, and productivity of honey bees. *Ukrainian Journal of Ecology*, 11(1), 344–349.
- Slivinska, L. G., Shcherbatyy, A. R., Lukashchuk, B. O., & Gutyj, B. V. (2020). The state of antioxidant protection system in cows under the influence of heavy metals. *Regulatory Mechanisms in Biosystems*, 11(2), 237–242.
- Slivinska, L. G., Shcherbatyy, A. R., Lukashchuk, B. O., Zinko, H. O., Gutyj, B. V., Lychuk, M. G., Chernushkin, B. O., Leno, M. I., Prystupa, O. I., Leskiv, K. Y., Slepokura, O. I., Sobolev, O. I., Shkromada, O. I., Kysterna, O. S., & Musienko, O. V. (2019). Correction of indicators of erythropoiesis and microelement blood levels in cows under conditions of technogenic pollution. *Ukrainian Journal of Ecology*, 9(2), 127–135.
- Slivinska, L. G., Vlizlo, V. V., Shcherbatyy, A. R., Lukashchuk, B. O., Gutyj, B. V., Drach, M. P., Lychuk, M. G., Maksymovych, I. A., Leno, M. I., Rusyn, V. I., Chernushkin, B. O., Fedorovych, V. L., Zinko, H. O., Prystupa, O. I., & Yaremchuk, V. Y. (2021). Influence of heavy metals on metabolic processes in cows. *Ukrainian Journal of Ecology*, 11(2), 284–291.
- Slivinska, L., Shcherbatyy, A., Gutyj, B., Lychuk, M., Fedorovych, V., Maksymovych, I., Rusyn, V., & Chernushkin, B. (2018). Parameters of erythropoiesis, acid resistance and population composition of erythrocytes of cows with chronic hematuria. *Ukrainian Journal of Ecology*, 8(1), 379–385.
- Stabler, D., Al-Esawy, M., Chennells, J. A., Perri, G., Robinson, A., & Wright, G. A. (2021). Regulation of dietary intake of protein and lipid by nurse-age adult worker honeybees. *Journal of Experimental Biology*, 224(3), 1–9.
- Takic, M., Zekovic, M., Terzic, B., Stojisavljevic, A., Mijuskovic, M., Radjen, S., & Ristic-Medic, D. (2021). Zinc Deficiency, Plasma Fatty Acid Profile and Desaturase Activities in Hemodialysis Patients: Is Supplementation Necessary? *Frontiers in Nutrition*, 8, 1–15.
- Tikhonova, O. M., Kyrylchuk, K. S., & Shapoval, V. P. (2020). Research of the gross content of nickel and arsenic in the diversion lanes of highways in the city of Sumy. *Bulletin of the Sumy National Agrarian University. Series "Agronomy and Biology"*, 2(40), 62–70.
- Trinkl, M., Kaluza, B. F., Wallace, H., Heard, T. A., Keller, A., & Leonhardt, S. D. (2020). Floral Species Richness Correlates with Changes in the Nutritional Quality of Larval Diets in a Stingless Bee. *Insects*, 11(125), 1–20.
- Vishechur, V. Y., Gutyj, B. V., Nischemenko, N. P., Kushnir, I. M., Salata, V. Z., Tarasenko, L. O., Khimych, M. S., Kushnir, V. I., Kalyn, B. M., Magrelo, N. V., Boiko, P. K., Kolotnytskyy, V. A., Velesyk, T., Pundyak, T. O., & Gubash, O. P. (2019). Effect of industry on the content of fatty acids in the tissues of the honey-bee head. *Ukrainian Journal of Ecology*, 9(3), 174–179.
- Vlizlo, V. V. (2012). *Laboratory research methods in biology, animal husbandry and veterinary medicine: a handbook*. Lviv, UA: Spolom Publishing House.
- Vozzhchova, R. A., Vlaschuk, A. M., & Drobit, O. S. (2021). Improvement of the ecological and remedial condition of soils on the basis of smart specialization. *Materials of the 10th All-Ukrainian scientific and practical conference of young scientists "Actual problems of agro-industrial production of Ukraine: sustainable development of agriculture in conditions of climate change"*. *Obroshyno*, Lviv-Obroshyno, 15–16.
- Wang, X., Zhong, Z., Chen, X., Hong, Z., Lin, W., Mu, X., Hu, X., & Zheng, H. (2021). High-Fat Diets with Differential Fatty Acids Induce Obesity and Perturb Gut Microbiota in Honey Bee. *International Journal of Molecular Sciences*, 22(2), 834–849.
- Yatsuk, I. P., & Balyuk, S. A. (2019). *Methodology of agrochemical certification of agricultural lands*. Kyiv (Leading normative document).
- Younus, H. (2018). Therapeutic potentials of superoxide dismutase. *International Journal of Health Sciences*, 12(3), 88–93.